

The end of 'health terrorism'? Investigating innovative approaches to substance abuse prevention

Clarissa Cook, University Department of Rural Health, University of Tasmania

There is a sense of urgency surrounding a key health problem of our time: high-risk drinking.¹ Misuse of alcohol is responsible for much of the acute and chronic disease burden, and is associated with mental health problems, suicides, and motor vehicle and other accidents.²⁻¹³ Risky drinking among young people, in particular, is widely regarded as an important public health issue not only because of the various harms incurred in the short term, but also because of the multitude of health, personal and social implications that are likely to affect people later in the life-course if such drinking patterns become entrenched.¹⁴ Australian youth in rural and remote communities are of particular concern since they consume alcohol at more harmful levels than their metropolitan counterparts.¹⁵

Despite substantial public investment and an array of different approaches, the 'problem' of binge-drinking has shown itself to be a highly complex and particularly intractable issue:

In our efforts to solve the problem of binge drinking, we have none of the precision that we like; it is not an infectious disease that can be controlled or eradicated by the application of so many units of some treatment, or prevented by the careful removal of clearly defined personal, social, or environmental factors that lead to illness.¹⁶

In Australia, as elsewhere, there is growing recognition that it is preferable to take a preventive approach to youth binge-drinking and alcohol problems more generally, rather than wait until the problem is apparent. Preventive programs are by no means a 'new invention', however – school-based alcohol abuse prevention programs have been part of Australian primary- and high-school education for many decades. Commentators have noted a number of phases of development in this country which have tended to mirror developments overseas.¹⁷

Early prevention work within schools tended to focus on the provision of information to students, particularly concerning the pharmacological dangers of substance use and the possible risky consequences of drinking. These programs often incorporated deliberate scare-tactics and have been labelled 'health terrorist'¹⁸ approaches due to the underlying assumption that scaring the living daylights out of people will 'scare the health into them'. Put simply, it was believed that "if young people just knew how horrible drugs were and what they did to their brains and bodies, then they would not use them".¹⁹ Sometimes more comprehensive school-based alcohol and drug education programs were delivered in conjunction with law enforcement agencies, with the aim of educating young people about the likely legal, social and health implications of the use of illicit drugs and the misuse of licit drugs.

Despite some residual elements of health terrorism within contemporary programs, the information approach as a stand-alone method of tackling high-risk drinking among youth was "an acknowledged failure by the late 1970s".²⁰ Ironically, some information-based programs have resulted in 'more educated drug users' as well as *increased* levels of use.¹⁹ The ensuing phase of school-based prevention took a more holistic approach – seeking to build the self-esteem of young people so that they were less vulnerable to the vagaries of substance abuse. Sometimes these programs included resistance training components that sought to 'innoculate' youth against overt peer-pressure to engage in risky behaviours. Over time such 'affective'

programs suffered the same fate as their predecessors the ‘information’ programs – they were gradually, if reluctantly recognised as having only limited efficacy.

With the exception of some more recent and more sophisticated ‘social influence’ programs²⁰, alcohol programs for young people have not achieved great success, either in Australia or elsewhere despite “good intentions and a parade of promising practices”.¹⁶ On the whole, alcohol educators here and overseas find themselves in a frustrating and disheartening position whereby, despite determined efforts, prevention programs generally fail to deliver sustained behavioural modification.¹⁷

LOOKING FOR ALTERNATIVES TO ‘HEALTH TERRORISM’

In searching for possible explanations for lack of effect it is necessary, we argue, to examine the assumptions underpinning the various prevention efforts. With respect to alcohol programs, information-based approaches assume that young people will be motivated to change by appeals to long-term health consequences or mortality. With respect to the so-called ‘affective’ and ‘inoculation’ approaches, there is an underlying assumption that low self-esteem is a significant causal factor in harmful patterns of alcohol consumption among young people. Similarly, although peer factors have repeatedly been shown to be fundamental to youth drinking behaviours²¹, it is conceivable that peer pressure does not operate in precisely the way program designers assumed that it does.

With such issues in mind, we argue in favour of the development of a ‘sociology of drinking’. We join d’Abbs²² in recognising that although the public health approach to alcohol-related problems is valuable from a descriptive and risk-factor identification perspective, it “fails to acknowledge the extent to which, and the many ways in which, drinking is a social as well as an individual act”. Furthermore, we insist that a sociological approach to alcohol consumption ‘matters very much’,

... not only because drinking is a social act, but because virtually the entire public health repertoire of policies and measures are... attempts to intervene in the social control of drinking.²²

As noted earlier, some of the more recent ‘social influence’ approaches to alcohol abuse prevention are yielding promising results. This could be because they incorporate environmental/cultural factors and acknowledge and utilise complex social control processes, rather than having a blinkered focus on the individual’s knowledge, values or personality in the manner characteristic of many ‘health terrorist’ approaches.

We argue that the pursuit of a theoretically sophisticated sociological approach to alcohol consumption represents an important way forward for rational program design and evidence-based policy development. One recent prevention approach that is gaining in popularity and deemed worthy of the label of ‘sociologically informed’, is known as ‘Social Norms’ (SN). SN has a theoretical basis in social-psychology, and draws upon theories of peer identity formation, conformity and cognitive dissonance.²³ A distinctive feature of SN is its clarification and utilisation of peer-related influences on behaviour. As explained by a pioneer of the approach,

Research has long pointed to the dramatic power of peer influence in adolescence and young adulthood, but what has not been adequately considered in previous research and prevention strategy is whether this peer influence comes simply from what other peers *actually* believe is the right thing to do and how they behave, or from what young people *think* their peers believe is right and how they *think* most others behave.²⁴

SN interventions do not seek to alter individuals' attitudes *per se*. Rather, they seek to understand the social processes that contribute to the creation and perpetuation of social environments that are supportive of risky alcohol consumption. Arguably, such an understanding is fundamental to any efforts to reduce alcohol-related harm by intervening in these processes.

The SN approach has been extensively employed in the United States, and has been heralded as an effective strategy for reducing alcohol-related harm in youthful populations by identifying and correcting such attitudinal and behavioural misperceptions. In the following section of this paper, we sketch out how the approach has developed since the foundational research was conducted nearly two decades ago, and consider whether or not the encouraging results achieved overseas would be likely to be achieved in the Australian context.

ABOUT THE SOCIAL NORMS APPROACH

The foundational research was undertaken in the late 1980s by social scientists Perkins and Berkowitz, who discovered widespread misperception of alcohol-related attitudes and behaviours among college students at Hobart and William Smith Colleges in upstate New York. Specifically, they found that students consistently overestimated how often and how much their peers drank, as well as overestimating their peers' support of risky drinking behaviours. Perkins and Berkowitz subsequently theorised that much high-risk activity stems from people wishing to, or feeling pressured to, conform to the behaviour and expectations of "imaginary peers".¹⁸

These early contentions have been supported by more recent studies – for instance, Beck and Trieman's finding that "teens' drinking behaviors are not driven so much by a need for peer approval or to be accepted by a group, but rather by *what is perceived of as normal behavior among one's close friends*".^{25,19} Essentially, what is problematic about misperception is the self-fulfilling prophecy²⁶ effect whereby the (often erroneous) assumption that 'everyone is doing it' leads to a situation where 'everyone does it'. Certainly, many studies demonstrate that perceptions of drinking norms predict, or are at least positively correlated with, individual drinking behaviours.^{21,27,28} However, just as inflated perceptions of drinking norms contribute to a social environment that is supportive of high-risk drinking, accurate norm perceptions will tend to have the opposite effect.¹⁷ Therein lies the 'secret weapon' of this important alternative to health terrorism:

The strategy of the social norms approach, put simply, is to communicate the truth about peer norms in terms of what the majority of students actually think and do, all on the basis of credible data drawn from the student population that is the target.²⁴

The basic stages of an SN intervention are as follows: The initial phase involves the collection of baseline self-report data about use and attitudes. These data are then analysed and the key messages are crafted, with an emphasis on positivity. (for example, "70% of Greentown High students have three or fewer drinks when they party"). Scare tactics and negative slants are notably absent. The next phase involves the incorporation of the key messages (i.e. the 'actual norms') into a media campaign utilising radio, flyers, screensavers, and newspaper ads, for example, that is then delivered intensively to the target population. The population from which the baseline data were collected is always the intended recipient of the media campaign, but sometimes additional groups (such as parents and teachers) are included. The media phase is then monitored for impact in terms of recognition and understanding of the message, changes to norm perceptions and resultant changes in behaviour.

Social norms interventions are rapidly gaining in popularity in the United States. In a survey of 4-year colleges nationwide in 1999, 20% of the colleges surveyed reported having conducted social norms marketing campaigns, and by 2001 this figure had risen to nearly 50%.²⁹ There is a growing body of evidence of encouraging and often dramatic reductions in high-risk drinking among target populations in metropolitan and non-metropolitan settings. For instance the University of Arizona reported a 29% reduction in 'heavy episodic drinking' over a three-year period.³⁰ Equivalent figures for other institutions include a 21% reduction over two years at the University of Missouri-Columbia, and a 44% reduction over 10 years at Northern Illinois University.³¹ Other institutions³² reported significant increases in the proportion of abstainers (teetotallers) among their student populations. Although the majority of SN interventions have been conducted at colleges and universities, the approach is also yielding promising results at high-schools.^{33,34}

Despite a growing band of enthusiastic followers, the SN approach does have its critics. Weschler, for example, recently argued that "...there is no evidence from scientifically rigorous evaluations supporting the effectiveness of...social norms marketing campaigns".²⁹ Although their conclusions have been refuted on methodological grounds³⁵, this group of Harvard-based academics remain vocal critics of the SN approach. Admittedly, there have been isolated examples of 'failed' SN interventions. Werch, for instance, reported that an intervention designed to prevent heavy episodic drinking among first-year college students "failed to produce any differences in self-reported alcohol use or alcohol-use risk indicators".^{36,37,38} However, the existence of such ineffective interventions do not, in themselves, constitute a satisfactory basis for dismissing the SN approach. We argue that the evidence base in support of the method is sufficiently large and robust to warrant detailed consideration of the potential 'fit' of SN within the Australian social, cultural and policy environments.

WOULD SOCIAL NORMS INTERVENTIONS BE LIKELY TO WORK IN AUSTRALIA?

Having learned something of the theoretical underpinnings of SN and the details of some interventions, we now face the task of considering whether or not the 'fit' between SN and the Australian policy and social environments is likely to be a comfortable one. Certainly, there are reasons to think that SN interventions might not be readily 'transplantable'. With few exceptions, virtually the entire body of evidence is US-based. There may be important cultural or social differences between Australia and the US (for instance, less pervasive peer orientation among adolescents) that would render SN interventions less effective in the former than in the latter. The American legal drinking age is 21 as opposed to 18, which might also have implications for program implementation.

Furthermore, the United States' "War on Drugs" is often held as the 'bastion of opposition' to Australia's drug policy position that is based on a 'harm reduction' approach.^{39,40} A detailed discussion of the similarities and differences between the drug policies of the two countries is not only outside the scope of this article, it is of limited value for the current discussion. What matters, we argue, is not how different the Australian and US drug policies are, but whether SN is itself compatible with a harm minimisation framework.

Although there has been some controversy surrounding the terms 'harm minimisation' and 'harm reduction'⁴¹ and the extent to which they are interchangeable, broadly speaking they refer to:

a policy of preventing the potential harms related to drug use rather than trying to prevent the drug use itself. Harm reduction accepts as a fact that drug use has persisted despite all efforts to prevent it and will continue to do so.⁴²

The principle of harm-minimisation/reduction provided the basis for Australia's National Campaign Against Drug Abuse (launched in 1985) as well as its successor, the National Drug Strategy.²² Critics of harm minimisation have suggested that it condones illicit drug use and other risky behaviours because it does not promote non-use, or even necessarily aim for a reduction in use. However, as Plant and his colleagues explain, harm minimisation is "neutral on the virtue or shame attached to such behaviours"⁴³ and although it does not seek to minimise alcohol intake *per se*, it is by no means incompatible with abstentionist aims.

There are good indications that SN interventions will fit comfortably within our harm minimisation policy framework. Unlike health promotion approaches that seek to scare people off behaviours because they are risky (or shame people out of them because they are 'bad'), SN approaches takes a neutral stance – they do not present alcohol consumption as either evil or virtuous. Importantly, there is an assumption that many young people do and will continue to consume alcohol – the challenge lies in finding evidence-based ways to diminish the likelihood of them harming either themselves or others in the process. SN is a promising candidate in this regard.

TRIALLING SOCIAL NORMS IN AUSTRALIA

We intend to conduct the first major Australian trial of the Social Norms approach to substance abuse prevention, and have recently submitted a funding application to the *Alcohol Education and Rehabilitation Foundation*. The project discussed below is dependent on a successful funding bid.

The proposed trial is a two year project (commencing January 2006) to implement SN interventions in two rural municipalities in Tasmania. It will aim to reduce binge drinking and alcohol-related harm among high-school aged youth in the Huon Valley and West Coast areas, focusing on the towns of Huonville, Geeveston, Rosebery and Queenstown. It will seek to identify technical procedures and practices, and community characteristics and conditions essential to the success of SN projects, with a key output of the project being the production of a resource kit on "How to conduct Social Norms Interventions". This kit will assist other Australian rural communities to conduct their own SN projects that 'replicate' the SN model but are still tailored to community needs and conditions.

The trial will take a collaborative, multi-disciplinary approach that acknowledges the importance of involving a diverse mix of individuals and institutions in prevention efforts.⁴⁴ This collaboration involves the:

- University of Tasmania
 - University Department of Rural Health (UDRH)
 - Tasmanian Institute of Law Enforcement Studies (TILES)
- Department of Police and Public Safety (Tasmania Police)
- Department of Health and Human Services (DHHS)
- Department of Education (DOE)
- two local high schools at each site
- local councils
- the business sector
- community representatives (e.g. youth).

The key members of the partnership all have substantial and well-established community linkages within the proposed trial sites as well as a history of successful collaborations with a variety of government and non-government organisations.

Students attending two public high schools in each area form the main target group. There are roughly twice as many students at the Huon Valley site as at the West Coast site (i.e. approximately 600 students at the former compared to only around 300 at the latter). Data on alcohol-related behaviours and attitudes will be collected from the students at three time points (baseline, mid-project and post-intervention) using an online survey. Data relating to alcohol-related harm among this group will also be collected from a variety of sources including police and health-services. The focus on youth in these particular age-groups is well-supported by the literature⁴⁵⁻⁴⁷, with strong agreement that the late primary/early high school years represent “the optimal time for initiating youth drug interventions” since it tends to coincide with the onset of experimentation.⁴⁸

Like many of the more recent SN interventions in the US, the Australian trial will take a broad community focus involving teachers and parents as well as students. Again, the inclusion of a parenting component in a youth-focused substance abuse prevention intervention is well supported by the literature.^{49,50} The trial will aim to identify and correct any misperceptions the parents might have of youth alcohol consumption in that community. An additional, though no less significant aim is to use the social norms approach to strengthen parenting behaviours that are supportive of safe alcohol consumption. Just like teens, parents’ behaviour can be influenced by erroneous perception of ‘peer’ (i.e. other parents’) behaviours and attitudes :

... if parents underestimate how frequently other parents are using certain protective strategies, this misperception may serve to undermine their own resolve to adopt those strategies or apply them consistently. Stated simply, it is harder for parents to uphold firm rules and standards when they believe they are among the few parents trying to do so.⁵¹

The broad, community-based approach of the proposed trial maximises potential reinforcement of the key messages.^{52,53} Furthermore, it seeks to prompt the ‘environmental’ level changes deemed necessary by Midford and colleagues, who argue that

curing or removing the individual problem drinker will not result in a reduction in alcohol-related harm, because the community dynamics which caused these problems are unchanged. In order to change the aggregate level of alcohol-related harm, environmental changes have to occur.⁴⁸

CONCLUSION

We are enthusiastic about the potential of the SN approach to reducing high-risk alcohol consumption among young people. It is an evidence-based prevention model that will hopefully avoid some of the ‘unintended consequences’ of media coverage and many of the standard scare-tactic health promotion approaches, which themselves contribute to the perception of the ‘normality’ of youth binge-drinking:

News accounts and other messages about student drinking that are designed to underscore the seriousness of the problem can have the unintended consequence of reinforcing the misperception that heavy drinking is the norm. Ironically, the very information that is designed to motivate corrective action may instead bolster a set of beliefs that make the problem more resistant to change.¹⁸

Although alcohol consumption has been the focus of most SN interventions in the US and will also be the focus of the Australian trial, the approach is by no means restricted to the area of substance abuse. There is a growing body of evidence that a variety of health and social justice

issues are amenable to change via the correction of misperceptions. For instance, encouraging results have been gained in relation to smoking^{51,54}, homophobic and racist behaviour^{55,56}, teenage pregnancy and sexual assault.^{57,58}

We are excited about conducting the first major Australian trial of the Social Norms approach, and are confident that our collaboration involving the University of Tasmania, the Tasmania Police, local and state government representatives, health care professionals, schools and rural community will work effectively towards achieving shared objectives. In the process of meeting important research priorities identified by the Australian government^{59,60} we will stimulate Australian debate about SN and provide evidence concerning its potential 'transplantation' to this country as a method for reducing alcohol-related harm. In adding to the body of knowledge about socio-cultural determinants of alcohol consumption, we will also contribute to the long-overdue development of a "Sociology of Drinking". We enthusiastically embrace this opportunity to examine an alternative to health terrorism that could revolutionise health promotion and make significant contributions to the health of rural and remote Australians.

REFERENCES

1. NHMRC. National Alcohol Guidelines: risks and benefits of consumption. Canberra: Commonwealth Department of Health and Aged Care; 2001.
2. Australian Institute of Health and Welfare. Drug Use in Australia and its Health Impact. In. Canberra: Australian Institute of Health and Welfare; 1999.
3. Baker S, B O'Neill B, Ginsburg M, Li M. The Injury Fact Book. New York: Oxford University Press; 1992.
4. Chikritzhs T, Catalano P, Stockwell T, Donath S, Ngo H, Young D, et al. Australian Alcohol Indicators 1990-2001. Perth: National Drug Research Institute, Curtin University of Technology and Turning Point Alcohol and Drug Centre; 2003.
5. Collins J, Messerschmidt P. Epidemiology of Alcohol-related Violence. Alcohol Health and Research World 1993;17(2):93-100.
6. d'Abbs P, Hunter E, Reser J, Martin D. Alcohol Related Violence in Aboriginal and Torres Strait Islander Communities: a literature review. Canberra: Australian Government Publishing Service; 1994. Report No.: 8.
7. Heale P, Chikritzhs T, Jonas H, Stockwell T, Dietze P. Estimated alcohol-caused deaths in Australia, 1990-1997. Drug and Alcohol Review 2002;21(2):121-129.
8. Jonas H, Dobson A, Brown W. Patterns of alcohol consumption in young Australian women: socio-demographic factors, health-related behaviour and physical health. Australian and New Zealand Journal of Public Health 2000;24(2):185-191.
9. Mason G, Wilson P. Alcohol and Crime. In. Canberra: Australian Institute of Criminology; 1989.
10. McBride N, Farrington F, Midford R. What harms do young Australians experience in alcohol use situations. Australian and New Zealand Journal of Public Health 2000;21(1):54-59.
11. Fombonne E. Suicidal behaviours in vulnerable adolescents: Timetrends and their correlates. British Journal of Psychiatry 1998(173):154-159.
12. Hall W, Farrell M. Co-morbidity between substance use and other mental disorders. Sydney: National Drug and Alcohol Research Centre; 1997.

13. White J, Humeniuk R. Alcohol Misuse and Violence: Exploring the Relationship. Adelaide: The Drug Offensive, National Symposium on Alcohol Misuse and Violence; 1994. Report No.: Report 2.
14. Loxley W, J Tombourou, T Stockwell, Haines B, Scott K, Godfrey C, Waters E, Patton G, Fordham R, Gray D, Marshall J, Ryder D, Sagggers S, Sanci L & Williams. The prevention of substance abuse, risk and harm in Australia: a review of the evidence: National Drug Research Institute; 2004.
15. Williams P. Alcohol-related Social Disorder and Rural Youth: Part 1 – Victims. Canberra: Australian Institute of Criminology; 1999 December.
16. Keeling R. The Political, Social and Public Health Problems of Binge Drinking in College. *Journal of American College Health* 2000;48(5):195-198.
17. Steffian G. Correction of Normative Misperceptions: An Alcohol Abuse Prevention Program. *Journal of Drug Education* 1999;29(2):115-138.
18. Wesley Perkins H, editor. *The Social Norms Approach to Preventing School and College Age Substance Abuse: A Handbook for Educators, Counsellors and Clinicians*. San Francisco: Jossey Bass; 2003.
19. Hogan J. *Substance Abuse Prevention : The Intersection of Science and Practice*. Boston: Allyn & Bacon; 2002.
20. Midford R, Munro G, McBride N, Ladzinski U. Principles that underpin effective school-based drug education. *Journal of Drug Education* 2002;32(4):363-386.
21. Borsari B, Carey K. Peer influences on college drinking: a review of the research. *Journal of Substance Abuse* 2001;13:391-424.
22. d'Abbs P. Silence of the sociologists: Indigenous alcohol use, harm minimisation and social control. *Health Sociology Review* 2002;10:33-52.
23. Perkins W. College student misperceptions of alcohol and other drug norms among peers: exploring causes, consequences, and implications for prevention programs. In: *Designing alcohol and other drug prevention programs in higher education: bringing theory into practice*. Newton, Mass: Higher Education Center for Alcohol and other Drug Prevention; 1997.
24. Perkins H, editor. *The Social Norms Approach to Preventing School and College Age Substance Abuse: A Handbook for Educators, Counsellors and Clinicians*. San Francisco: Jossey Bass; 2003.
25. Beck K, Treiman K. The Relationship of Social Context of Drinking, Perceived Social Norms, and Parental Influence to Various Drinking Patterns of Adolescents. *Addictive Behaviors* 1996;21(5):633-644.
26. Merton R. The Self-Fulfilling Prophecy. In: *Social Theory and Social Structure*. New York: Free Press; 1957.
27. Thombs D, B Wolcott and L Farkash. Social Context, Perceived Norm and Drinking Behavior in Young People. *Journal of Substance Abuse* 1997;9:257-267.
28. Page R, Scanlan A, Gilbert L. Relationship of the estimation of binge drinking among college students and personal participation in binge drinking: Implications for health education and promotion. *Journal of Health Education* 1999;30:98-103.
29. Weschler Hea. Colleges Respond to Student Binge Drinking: Reducing Student Demand or Limiting Access. *Journal of American College Health* 2004;52(4):159-168.
30. Glider P, Midyett S, Mills-Novoa B, Johannessen K, Collins C. Challenging the collegiate rite of passage: a campus-wide social marketing media campaign to reduce binge-drinking. *Journal of Drug Education* 2001;31(207-220).

31. Haines MP. A social norms approach to preventing binge drinking at colleges and universities. Newton, MA: Higher Education Centre for Alcohol and other Drug Prevention, Education Development Center Inc; 1996.
32. Peeler M, Far J, Miller J, Brigham T. An analysis of the effects of a program to reduce heavy drinking among college students. *Journal of Alcohol and Drug Education* 2000;45:39-54.
33. Linkenbach J. Imaginary peers and the reign of error. *Prevention Connection* 1999;3:1-5.
34. Johannessen K, Collins C, Mills-Novoa B, Glider P. A Practical Guide to Alcohol Abuse Prevention: A Campus Case Study in implementing social norms and environmental management approaches. Tucson, Arizona: Campus Health Service, University of Arizona; 1999.
35. Perkins H, Linkenbach J. Harvard Study of Social Norms Deserves "F" Grade for Flawed Research Design. In; 2003.
36. Werch C. Results of a social norm intervention to prevent binge-drinking among first-year residential college students. *Journal of American College Health* 2000;49:85-92.
37. Clapp J, Lange J, Russe C, Shillington A, Voas R. A failed social marketing campaign. *Journal of Studies on Alcohol* 2003;64:409-414.
38. Trockel M, S Williams and J Reis. Considerations for more effective Social Norms Based Alcohol Education on Campus: An Analysis of Different Theoretical Conceptualizations in Predicting Drinking Among Fraternity Men. *Journal of Studies on Alcohol* 2003;64:50-59.
39. Roche A, Evans K, Stanton W. Harm reduction: roads less travelled to the Holy Grail. *Addiction* 1997;92(9):1207-1212.
40. Wink W. Getting off drugs: the legalisation option. *Friends Journal* 1996(February).
41. Single E, Rohl T. The National Drug Strategy: Mapping the future. An evaluation of the National Drug Strategy 1993-1997: A report commissioned by the Ministerial Council on Drug Strategy. Canberra: Australian Government Publishing Service; 1997.
42. Duncan D, Nicholson T, Clifford P, Hawkins W, Petosa R. Harm reduction: an emerging new paradigm for drug education. *Journal of Drug Education* 1994;24(4):281-290.
43. Plant M, Single E, Stockwell T. Introduction: Harm minimisation and alcohol. In: Plant M, Single E, Stockwell T, editors. *Alcohol: Minimising the Harm*. London: Freedom Association Books; 1997.
44. Roche A, Stockwell T. Putting prevention back on the agenda. *Drug and Alcohol Review* 2004;23(1):3-4.
45. Johnston L, O' Malley P, Bachman J. *Drug Use, Drinking and Smoking: National Survey Results from High School, College and Young Adult Populations, 1975-1988*. Washington, DC: National Institute on Drug Abuse; 1989. Report No.: DHHS Pub. No. (ADM) 89-1638.
46. Dielman T. School-based research on the prevention of adolescent alcohol use and misuse: Methodological issues and advances. *Journal of Research on Adolescence* 1994;4(2):271-293.
47. Duncan T, Duncan S, Hops H. The effects of family cohesiveness and peer encouragement on the development of adolescent alcohol use: A cohort-sequential approach to the analysis of longitudinal data. *Journal of Studies on Alcohol* 1994;55:588-599.
48. Midford R, Stockwell T, Gray D. Prevention of alcohol-related harm: community-based interventions. In: Australia Co, editor. *National Alcohol Research Agenda: A Supporting Paper to the National Alcohol Strategy A Plan for Action 2001 to 2003-4*. Canberra: Publications Production Unit, Commonwealth Department of Health and Ageing; 2002. p. 91-99.

49. Rohrbach L, Hodgson C, Broder B, Montgomery S, Flay B, Hansen W, et al. Parental participation in drug use prevention: Results from the Midwestern Prevention Project. *Journal of Research on Adolescence* 1994;4(2):295-317.
50. Beck K, Lockhart S. A model of parental involvement in adolescent drinking and driving. *Journal of Youth and Adolescence* 1992;21(1):35-51.
51. Hancock H, Henry N. Perceptions, norms and tobacco use in college residence hall freshmen: evaluation of a social norms marketing intervention. In: Perkins HW, editor. *The Social Norms Approach to Preventing School and College Age Substance Abuse: A Handbook for Educators, Counsellors and Clinicians*. San Francisco: Jossey Bass; 2003. p. 247-258.
52. Perry C, Kelder S. Prevention. *Annual Review of Addictions Research and Treatment* 1992:453-472.
53. Perry C, Williams C, Veblen-Mortenson S, Toomey T, Komro K, Anstine P, et al. Project Northland: Outcomes of a community-wide alcohol use prevention program during early adolescence. *American Journal of Public Health* 1996;86(7):956-965.
54. Linkenbach J, Perkins HW. MOST of Us are Tobacco Free: An eight-month social norms marketing campaign reducing youth initiation of smoking in Montana. In: Perkins HW, editor. *The Social Norms Approach to Preventing School and College Age Substance Abuse: A Handbook for Educators, Counsellors and Clinicians*. San Francisco: Jossey Bass; 2003. p. 247-258.
55. Fabiano P. Using a Social Norms approach for building just and non-violent communities. In: *Third Annual Conference on Social Norms: Science-based prevention from theory to practice*; 2000 July 26-28; Denver, Colorado; 2000.
56. Smolinsky T. What do we really think?: A group exercise to increase heterosexual ally behavior. In: *The Report on Social Norms*. Little Falls, NJ: Paperclip Communications; 2002.
57. Berkowitz A. Fostering Men's Responsibility for Preventing Sexual Assault. In: Schewe P, editor. *Preventing Violence in Relationships: Interventions Across the Life Span*. Washington, D.C. ?; 2002.
58. Bruce S. The "A Man" campaign: Marketing social norms to men to prevent sexual assault. In: *The Report on Social Norms*. Little Falls, NJ: PaperClip Communications; 2002.
59. Commonwealth of Australia. *National Alcohol Research Agenda: A Supporting Paper to the National Alcohol Strategy A Plan for Action 2001 to 2003-4*. Canberra: Publications Production Unit, Commonwealth Department of Health and Ageing; 2002 March.
60. Roche A, Stockwell T. Prevention of alcohol-related harm: public policy and health. In: Australia Co, editor. *National Alcohol Research Agenda: A Supporting Paper to the National Alcohol Strategy A Plan for Action 2001 to 2003-4*. Canberra: Publications Production Unit, Commonwealth Department of Health and Ageing; 2002. p. 57-73.

PRESENTER

Clarissa Cook is a Research Fellow at the University Department of Rural Health at the University of Tasmania. She delivers research training to primary health professionals as part of the PHCRED initiative, and frequently advises state government employees on research and evaluation design. Clarissa seeks to emphasise and support theoretically informed practice, multi-disciplinary and collaborative undertakings, innovative service delivery, and rural community engagement.